

Osteoarthritis and Cartilage



Corrigendum

Corrigendum to “The relationship between fibrogenic TGFβ1 signaling in the joint and cartilage degradation in post-injury osteoarthritis” [Osteoarthritis and Cartilage 19 (2011) 1081–1090]



A. Plaas ^{†‡}, J. Velasco [‡], D.J. Gorski [‡], J. Li [†], A. Cole [‡], K. Christopherson ^{‡§}, J.D. Sandy ^{‡||} *

[†] Department of Internal Medicine (Rheumatology), Rush University Medical Center, Chicago, IL, USA

[‡] Department of Biochemistry, Rush University Medical Center, Chicago, IL, USA

[§] Department of Internal Medicine (Hematology/Cell Therapy), Rush University Medical Center, Chicago, IL, USA

^{||} Department of Orthopedic Surgery, Rush University Medical Center, Chicago, IL, USA

The authors regret that references 137–180 were missing from the original article published. This caused citations within [Table 1](#) to not be referenced and so [Table 1](#) has been corrected and appears below, along with the missing references.

The authors would like to apologise for any inconvenience caused.

Table 1
Wound healing mediators in dermal and joint tissues showing their source and major effects on cellular responses

Factor	Joint tissue source	Joint tissue repair responses	Wound cell source [20–23]	Wound healing responses [20–23]
EGF	SY[133]; OA SF[134]	Chondrocyte proliferation; ion transport, decreased matrix production	P,M,F	Epithelialization
FGF-2	SY[135–137]; AT[138]; OP[137]; OA SF[139–142]	Anti-apoptotic; prochondrogenic	M,EP,END,F	Angiogenesis Granulation tissue ECM production
TGFβ1	CA[143]; SY[144]; OP[137, 145]; OA SF[47, 48]	Pro-catabolic (MMP-13); chondrocyte hypertrophy	P,M,EP,END,F	Epithelialization, Granulation Tissue Fibroplasia
BMPs	SY[152]; CA[146–148]; BO[149, 150]; OA SF[151]	Prochondrogenic; Osteophytes	SC	Hairfollicle formation
PDGF	CA[143]; SY[153, 154]; OA SF[47]	Stimulates reparative responses in fibrochondrocytes; anti-hypertrophic	P,M,F	Granulation tissue Fibroplasia Contraction
VEGF	CA[155–159,163]; SY[158–161]; AT[138]; OA SF[47, 162]	Delays reparative responses in meniscus and CA	P,N,M,END,F	Angiogenesis
IL1β	CA[143]; SY[180]; post ACLT SF[164–167]; OA SF[48, 167–170]	CA and meniscal matrix degradation	N,M,EP	Inflammation
IL6	CA[171, 172]; AT[138, 173]; PC[174]; Post ACL SF[164, 175]; OA SF[48, 169]	CA matrix degradation	N,M,EP	Epithelialization
TNFα	CA[143]; SY[176]; AT[138]; Post ACLT SF[167,170,175, 177,178]; OA SF[48,168,169].	CA matrix degradation	N,M,EP	Inflammation Epithelialization

Abbreviations: ACLT: Anterior cruciate ligament tear; AT: Adipose tissue; OP: Osteophyte; PC: Plasma Cells; END: endothelial cells; EP: epithelial cells; F: Fibroblasts; M: Macrophages; N: Neutrophils; P: Platelets.

Missing References

137. Hosaka S, Shah MR, Barquin N, Haines GK, Koch AE. Expression of basic fibroblast growth factor and angiogenin in arthritis. *Pathobiology* 1995;63:249–56.
138. Uchino M, Izumi T, Tominaga T, Wakita R, Minehara H, Sekiguchi M, *et al.* Growth factor expression in the osteophytes of the human femoral head in osteoarthritis. *Clin Orthop Relat Res* 2000;119–25.
139. Ushiyama T, Chano T, Inoue K, Matsusue Y. Cytokine production in the infrapatellar fat pad: another source of cytokines in knee synovial fluids. *Ann Rheum Dis* 2003;62:108–12.
140. Gudbjornsson B, Christofferson R, Larsson A. Synovial concentrations of the angiogenic peptides bFGF and VEGF do not discriminate rheumatoid arthritis from other forms of inflammatory arthritis. *Scand J Clin Lab Invest* 2004;64:9–15.

DOI of original article: <http://dx.doi.org/10.1016/j.joca.2011.05.003>.

* Address correspondence and reprint requests to: J.D. Sandy, Department of Biochemistry, Rush University Medical Center, Chicago, IL, USA.

E-mail address: jsandy44@gmail.com (J.D. Sandy).

<http://dx.doi.org/10.1016/j.joca.2014.05.002>

1063-4584/© 2014 Osteoarthritis Research Society International. Published by Elsevier Ltd. All rights reserved.

141. Sato J, Segami N, Kaneyama K, Konishi H, Yoshitake Y, Nishikawa K. Levels of fibroblast growth factor 2 in synovial fluids in human patients with internal derangement of the temporomandibular joint. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;96:673–79.
142. Orito K, Koshino T, Saito T. Fibroblast growth factor 2 in synovial fluid from an osteoarthritic knee with cartilage regeneration. *J Orthop Sci* 2003;8:294–300.
143. Schmal H, Mehlhorn A, Stoffel F, Kostler W, Sudkamp NP, Niemeyer P. In vivo quantification of intraarticular cytokines in knees during natural and surgically induced cartilage repair. *Cytotherapy* 2009;11:1065–75.
144. Moos V, Fickert S, Muller B, Weber U, Sieper J. Immunohistological analysis of cytokine expression in human osteoarthritic and healthy cartilage. *J Rheumatol* 1999;26:870–9.
145. Yoshida H, Yoshida T, Iizuka T, Sakakura T, Fujita S. The expression of transforming growth factor beta (TGF-beta) in the synovial membrane of human temporomandibular joint with internal derangement: a comparison with tenascin expression. *J Oral Rehabil* 1999;26:814–20.
146. Verdier MP, Seite S, Guntzer K, Pujol JP, Boumediene K. Immunohistochemical analysis of transforming growth factor beta isoforms and their receptors in human cartilage from normal and osteoarthritic femoral heads. *Rheumatol Int* 2005;25:118–24.
147. Kwong FN, Hoyland JA, Freemont AJ, Evans CH. Altered relative expression of BMPs and BMP inhibitors in cartilaginous areas of human fractures progressing towards nonunion. *J Orthop Res* 2009;27:752–57.
148. Bobinac D, Spanjol J, Marinovic M, Zoricic Cvek S, Maric I, Cicvaric T, et al. Expression of bone morphogenetic proteins, cartilage-derived morphogenetic proteins and related receptors in normal and osteoarthritic human articular cartilage. *Coll Antropol* 2008;32(Suppl 2):83–7.
149. Chen AL, Fang C, Liu C, Leslie MP, Chang E, Di Cesare PE. Expression of bone morphogenetic proteins, receptors, and tissue inhibitors in human fetal, adult, and osteoarthritic articular cartilage. *J Orthop Res* 2004;22:1188–92.
150. Zoricic S, Maric I, Bobinac D, Vukicevic S. Expression of bone morphogenetic proteins and cartilage-derived morphogenetic proteins during osteophyte formation in humans. *J Anat* 2003;202:269–77.
151. Nauth A, Ristiniemi J, McKee MD, Schemitsch EH. Bone morphogenetic proteins in open fractures: past, present, and future. *Injury* 2009;40(Suppl 3):S27–31.
152. Chubinskaya S, Frank BS, Michalska M, Kumar B, Merrihew CA, Thonar EJ, et al. Osteogenic protein 1 in synovial fluid from patients with rheumatoid arthritis or osteoarthritis: relationship with disease and levels of hyaluronan and antigenic keratan sulfate. *Arthritis Res Ther* 2006;8:R73.
153. Schmal H, Niemeyer P, Zwingmann J, Stoffel F, Sudkamp NP, Mehlhorn AT. Association between expression of the bone morphogenetic proteins 2 and 7 in the repair of circumscribed cartilage lesions with clinical outcome. *BMC Musculoskelet Disord* 2010;11:170.
154. Pohlars D, Huber R, Ukena B, Kinne RW. Expression of platelet-derived growth factors C and D in the synovial membrane of patients with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum* 2006;54:788–94.
155. Remmers EF, Sano H, Wilder RL. Platelet-derived growth factors and heparin-binding (fibroblast) growth factors in the synovial tissue pathology of rheumatoid arthritis. *Semin Arthritis Rheum* 1991;21:191–99.
156. Lingaraj K, Poh CK, Wang W. Vascular endothelial growth factor (VEGF) is expressed during articular cartilage growth and re-expressed in osteoarthritis. *Ann Acad Med Singapore* 2010;39:399–403.
157. Franses RE, McWilliams DF, Mapp PI, Walsh DA. Osteochondral angiogenesis and increased protease inhibitor expression in OA. *Osteoarthritis Cartilage* 2010;18:563–71.
158. Murata M, Yudoh K, Masuko K. The potential role of vascular endothelial growth factor (VEGF) in cartilage: how the angiogenic factor could be involved in the pathogenesis of osteoarthritis? *Osteoarthritis Cartilage* 2008;16:279–86.
159. Enomoto H, Inoki I, Komiya K, Shiomi T, Ikeda E, Obata K, et al. Vascular endothelial growth factor isoforms and their receptors are expressed in human osteoarthritic cartilage. *Am J Pathol* 2003;162:171–81.
160. Sato J, Segami N, Nishimura M, Kaneyama K, Demura N, Yoshimura H. Relation between the expression of vascular endothelial growth factor in synovial tissues and the extent of joint effusion seen on magnetic resonance imaging in patients with internal derangement of the temporomandibular joint. *Br J Oral Maxillofac Surg* 2003;41:88–94.
161. Haywood L, McWilliams DF, Pearson CI, Gill SE, Ganesan A, Wilson D, et al. Inflammation and angiogenesis in osteoarthritis. *Arthritis Rheum* 2003;48:2173–77.
162. Sato J, Segami N, Yoshitake Y, Nishikawa K. Correlations of the expression of fibroblast growth factor-2, vascular endothelial growth factor, and their receptors with angiogenesis in synovial tissues from patients with internal derangement of the temporomandibular joint. *J Dent Res* 2003;82:272–77.
163. Kim KS, Choi HM, Lee YA, Choi IA, Lee SH, Hong SJ, et al. Expression levels and association of gelatinases MMP-2 and MMP-9 and collagenases MMP-1 and MMP-13 with VEGF in synovial fluid of patients with arthritis. *Rheumatol Int* 2010.
164. Matsumoto K, Honda K, Ohshima M, Yamaguchi Y, Nakajima I, Micke P, et al. Cytokine profile in synovial fluid from patients with internal derangement of the temporomandibular joint: a preliminary study. *Dentomaxillofac Radiol* 2006;35:432–41.
165. Cameron ML, Fu FH, Paessler HH, Schneider M, Evans CH. Synovial fluid cytokine concentrations as possible prognostic indicators in the ACL-deficient knee. *Knee Surg Sports Traumatol Arthrosc* 1994;2:38–44.
166. Irie K, Uchiyama E, Iwaso H. Intraarticular inflammatory cytokines in acute anterior cruciate ligament injured knee. *Knee* 2003;10:93–6.
167. Catterall JB, Stabler TV, Flannery CR, Kraus VB. Changes in serum and synovial fluid biomarkers after acute injury (NCT00332254). *Arthritis Res Ther* 2010;12:R229.
168. Zysk SP, Fraunberger P, Veihelmann A, Dorger M, Kalteis T, Maier M, et al. Tunnel enlargement and changes in synovial fluid cytokine profile following anterior cruciate ligament reconstruction with patellar tendon and hamstring tendon autografts. *Knee Surg Sports Traumatol Arthrosc* 2004;12:98–103.
169. Westacott CI, Whicher JT, Barnes IC, Thompson D, Swan AJ, Dieppe PA. Synovial fluid concentration of five different cytokines in rheumatic diseases. *Ann Rheum Dis* 1990;49:676–81.
170. Holt I, Cooper RG, Denton J, Meager A, Hopkins SJ. Cytokine inter-relationships and their association with disease activity in arthritis. *Br J Rheumatol* 1992;31:725–33.
171. Marks PH, Donaldson ML. Inflammatory cytokine profiles associated with chondral damage in the anterior cruciate ligament-deficient knee. *Arthroscopy* 2005;21:1342–47.

172. Moktar NM, Yusof HM, Yahaya NH, Muhamad R, Das S. The transcript level of interleukin-6 in the cartilage of idiopathic osteoarthritis of knee. *Clin Ter* 2010;161:25–8.
173. Kaneyama K, Segami N, Sato J, Nishimura M, Yoshimura H. Interleukin-6 family of cytokines as biochemical markers of osseous changes in the temporomandibular joint disorders. *Br J Oral Maxillofac Surg* 2004;42:246–50.
174. Distel E, Cadoudal T, Durant S, Poignard A, Chevalier X, Benelli C. The infrapatellar fat pad in knee osteoarthritis: an important source of interleukin-6 and its soluble receptor. *Arthritis Rheum* 2009;60:3374–77.
175. Doss F, Menard J, Hauschild M, Kreutzer HJ, Mittlmeier T, Muller-Steinhardt M, *et al.* Elevated IL-6 levels in the synovial fluid of osteoarthritis patients stem from plasma cells. *Scand J Rheumatol* 2007;36:136–9.
176. Cameron M, Buchgraber A, Passler H, Vogt M, Thonar E, Fu F, *et al.* The natural history of the anterior cruciate ligament-deficient knee. Changes in synovial fluid cytokine and keratan sulfate concentrations. *Am J Sports Med* 1997;25:751–54.
177. Wassilew GI, Lehnigk U, Duda GN, Taylor WR, Matziolis G, Dinybil C. The expression of proinflammatory cytokines and matrix metalloproteinases in the synovial membranes of patients with osteoarthritis compared with traumatic knee disorders. *Arthroscopy* 2010;26:1096–104.
178. Higuchi H, Shirakura K, Kimura M, Terauchi M, Shinozaki T, Watanabe H, *et al.* Changes in biochemical parameters after anterior cruciate ligament injury. *Int Orthop* 2006;30:43–7.
179. Elsaid KA, Fleming BC, Oksendahl HL, Machan JT, Fadale PD, Hulstyn MJ, *et al.* Decreased lubricin concentrations and markers of joint inflammation in the synovial fluid of patients with anterior cruciate ligament injury. *Arthritis Rheum* 2008;58:1707–15.
180. Pelletier JP, McCollum R, Cloutier JM, Martel-Pelletier. Synthesis of metalloproteinases and IL-6 in human osteoarthritic synovial membrane is an IL-1 mediated process. *J Rheumatol Suppl* 1995;43:109–14.